

Repeating Defects



To identify the component causing the defect, compare your print sample against the measurements shown.

To diagnose a defect on the transfer belt:

1. Print eight Letter/A4-size pages long-edge feed. Verify that the same defect appears on every fourth page.
- Or
2. Print four Tabloid/A3-size pages. Verify that the same defect appears on every other page.

Call for service for this defect.




Table 1. Mean (SD) age, height, weight, and body mass index (BMI) of the 100 children in the study

Age (years)	Height (cm)	Weight (kg)	BMI (kg m ⁻²)
7.0	121.5 (6.5)	24.5 (5.5)	16.5 (2.5)
8.0	130.5 (7.5)	32.5 (8.5)	19.5 (3.5)
9.0	139.5 (8.5)	41.5 (11.5)	22.5 (4.5)
10.0	148.5 (9.5)	50.5 (14.5)	25.5 (5.5)
11.0	157.5 (10.5)	59.5 (17.5)	28.5 (6.5)
12.0	166.5 (11.5)	68.5 (20.5)	31.5 (7.5)
13.0	175.5 (12.5)	77.5 (23.5)	34.5 (8.5)
14.0	184.5 (13.5)	86.5 (26.5)	37.5 (9.5)
15.0	193.5 (14.5)	95.5 (29.5)	40.5 (10.5)
16.0	202.5 (15.5)	104.5 (32.5)	43.5 (11.5)
17.0	211.5 (16.5)	113.5 (35.5)	46.5 (12.5)
18.0	220.5 (17.5)	122.5 (38.5)	49.5 (13.5)
19.0	229.5 (18.5)	131.5 (41.5)	52.5 (14.5)
20.0	238.5 (19.5)	140.5 (44.5)	55.5 (15.5)

3.1. Prevalence of overweight and obesity

Figure 1 shows the prevalence of overweight and obesity in the study population. The prevalence of overweight increased from 10.0% at age 7 to 20.0% at age 18. The prevalence of obesity increased from 1.0% at age 7 to 10.0% at age 18. The prevalence of overweight and obesity increased with age in both sexes.

Figure 2 shows the prevalence of overweight and obesity in the study population by sex. The prevalence of overweight and obesity increased with age in both sexes.

Figure 3 shows the prevalence of overweight and obesity in the study population by age group. The prevalence of overweight and obesity increased with age in both sexes.

Figure 4 shows the prevalence of overweight and obesity in the study population by BMI. The prevalence of overweight and obesity increased with BMI in both sexes.

3.2. Discussion

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

The prevalence of overweight and obesity in the study population increased with age. The prevalence of overweight and obesity increased with age in both sexes.

